

Carry your treatment details

Print, cut out and fold. Follow your treating hospital's contact plan.

SACTCHECK · TREATMENT CARD

Adjuvant abemaciclib + endocrine therapy

NCCP 00619 v4a

The usual abemaciclib starting dose is 150 mg twice daily; a lower dose or altered schedule may be prescribed. Abemaciclib is taken twice daily continuously for up to two years with endocrine therapy, unless recurrence or unacceptable toxicity requires an earlier change. A four-week dispensing block is not a rest cycle. Endocrine treatment normally continues beyond abemaciclib.



Prescriber / next review: _____

Contact plan

Oncology team: Insert numbers here · Use your hospital's opening hours.

Out of hours: Insert numbers here.

Evenings / overnight: agreed out-of-hours service or emergency department.

Emergency: 112 / 999.

Own hospital / contact: _____

Report new breathlessness, chest symptoms, severe diarrhoea, inability to drink, jaundice, a painful swollen leg or a severe rash promptly. You do not need to identify which medicine caused the symptom.

<https://sactcheck.com/patient/00619/> · no patient information encoded.

SACTCheck · Developed by Dr Paul O'Brien · v0.84.21 · Sources checked 2026-10-25

Source reconciled; independent clinical, pharmacy and patient/PPI review pending